


Translating Research to Public Health Action: Asthma, Pest Infestation, and Unsafe Pesticide Use In New York City

Thomas Matte
Dan Kass
New York City Department of Health and Mental Hygiene

Overview

- Pediatric asthma - New York City context and public health programs
- DOHMH IPM, pesticide use reduction, and surveillance initiatives
- Public health agency-academic collaboration

Asthma hospitalization rates among children have fallen, but large disparities remain.



Health care and environmental context

- Use of controller medications increasing
- ICS under used in selected populations studied
- NYC MSA in non-attainment status for ozone and PM 2.5
- Insufficient data on neighborhood variation in ambient air pollution
- Asthma disparities are mirrored by disparities in pest infestation and unsafe pesticide use

NYC Asthma Initiative Goals

- Improving asthma clinical management
 - Severity assessment
 - Appropriate medication use
- Promoting asthma self-management skills
- Improving care coordination
- Reducing asthma trigger exposure

NYC Asthma Initiative Programs

Programs	Approach
Managing Asthma in Schools, daycares	Care coordination; self-management education
Community asthma care coordinators	Care coordination
Training institute, public health detailing	Guideline dissemination/promotion
NYC Asthma Partnership	Coalition building, policy development
Community IPM	Allergen reduction in homes


Other local initiatives with asthma control benefits

- **Tobacco control** – bar and restaurant smoking ban, increased cigarette taxes, large-scale free NRT distribution
- **Reducing diesel particulate emissions** in MTA fleet, construction equipment, and school busses
- NYC HANES, CHS, Automated Student Health Record, Primary Care Information Project - **Enhanced surveillance of asthma and relevant exposures**

Home allergen reduction for inner- city children with asthma

- Controlled studies - variations on home education/ coaching, provision of tools/ supplies, professional extermination, and repeated visits.
- Cost ~ \$1000-\$2000 per home
- Reduction in allergen levels
- Fewer symptom days, in some cases urgent health care use

Sources: Kreiger et al. AJPH 2005;95:652-659; Carter et al. JACI 2001;108:732-7; Morgan et al. NEJM 2004;351:1068-80; Eggleston et al. AAAI;2005;95:496-7.



Collaboration: DOHMH and the Columbia CCEH

- NYC Public Housing Authority IPM intervention design and evaluation
- Pesticide exposure surveillance
- Policy formation and implementation

Public Health Rationale for Promoting IPM

- Targets a contributor to asthma disparities
- IPM overlaps with interventions that address other housing-related health concerns
- Health risks from residential pesticide use
 - ~1,000 annual referrals to Poison Control Center and > 60 hospitalizations
 - Subclinical developmental toxicity

Why start with NYC Housing Authority?

- Largest PHA in North America
 - 179,000 apartments, 344 developments, 2,700 buildings
 - ~ 9% of NYC's rental apartments, 5% of population
 - 75,000 families, 413,000 authorized residents
- Developments concentrated in areas with high asthma prevalence and morbidity
- Likelihood of IPM success and sustainability: quality low-income housing and centralized pest control policies

Intervention Design and Evaluation

- In 2001, DOHMH expanded pilot IPM effort with NYC Housing Authority
- Goal: to test IPM versus traditional pest control for impact on pest burden in low income housing.
- DOHMH partnered with David Evans and the CCCEH to expand evaluation to include impact on allergens and, on exploratory basis, asthma symptoms.
- CCEHC provided multi-disciplinary expertise in design, statistical analysis, and asthma evaluation as well as important resource personnel.

NYCHA's IPM Intervention


The IPM intervention involves:

- Professional cleaning to remove food sources and cockroach frass in **kitchen and bathrooms**.
- Caulking and sealing kitchen cabinets and other points of entry.
- Use of MaxForce gels & bait stations, and boric acid powder.
- Residents receive a garbage can with a lid, food storage containers, and cleaning supplies.



Evaluation Design

Intervention Buildings



Control Buildings



Data collected during each home visit:

- Questionnaire about pest sightings, pesticide use, asthma symptoms, and opinions of building conditions
- Cockroach and mouse populations monitored for 1 week
- Dust sample from kitchen floor and up to 3 beds

Results at 6 Months IPM Units vs. Control Units

- Cockroach counts reduced > 80%
- Rodent sightings reduced > 50%
- Cockroach allergen reduced in kitchens and bedrooms

IPM Evaluation Impact

- Largest evaluation to date of public housing IPM intervention
- NYCHA implementation:
 - Established permanent IPM team hired from residents
 - Suspended the use of residual spray pesticides and pyrethroids for routine pest control
 - Expanding IPM to additional developments
- Manuscript in preparation

Pesticide exposure surveillance

- NYC Health and Nutrition Examination Survey (HANES) - Pesticide Biomonitoring
 - Adopted Whyatt, Barr et al. methods for exposure monitoring of pyrethroids and organophosphates
 - Columbia shared pre-publication results of their cohort studies to help determine biomonitoring priorities
- CCCEH serves on NYC Environmental Public Health Tracking Pesticide Surveillance Advisory Committee

Reducing Unsafe Pesticide Use

- CCCEH – Expert testimony on proposed City Council legislation to prohibit certain pesticide use on City-owned property
- Local Law 37 Features
 - Prohibits use of highly toxic pesticides, products containing carcinogens and reproductive/developmental hazards
 - Commits NYC to adopt IPM for its own property
 - Enhances surveillance of pesticide use, Potential source of future data for studies and collaboration.

Scientific Collaboration

- Pesticide Use – Exposure Validation Study
 - Background
 - New York State Pesticide Use Registry records address-application-product specific data
 - DOHMH is studying relationship between birth outcomes and commercial pesticide use
 - Wyatt et al. following mother-child pairs prospectively for pesticide exposure and outcomes
 - Opportunity to validate pesticide use data with address-specific exposure data, linked at address and time.

Translating research to public health interventions

- Finding the right balance of reach and effectiveness
- Credible health impact projections for setting priorities and stakeholder support
- Availability of usable local surveillance data
- Policy, institutional, structural changes to achieve broad reach and sustainability – e.g. school health MAF change
- Interagency collaboration
- Academic collaboration to build needed capacity